

FLOWERING PHENOLOGY AND FERTILITY OF SOUR CHERRY (*Prunus cerasus* L.) CULTIVARS SELECTED IN HUNGARY

József Nyéki¹, Tibor Szabó², Zoltán Szabó¹

¹University of Debrecen, Centre of Agricultural Sciences,
H-4032 Debrecen, Böszörményi út 138., Hungary,
Phone: (36) 52-508-477, e-mail: erdodine@helios.date.hu

²Research and Extension Centre for Fruit Growing, H-4244 Újfehértó, Vadastag 2.
Phone: (36) 42-290-822, Fax: (36) 42-290-822

S u m m a r y

Experiments were conducted between 1972 and 2002 at three sites in Hungary. At Érd 97, Helvécia 10, and Újfehértó 3 cultivars of sour cherry were studied in variety collections. Observations were made on the flowering phenology (start, main time, end and length of the bloom period), on the flowering dynamics (the rate of the open flowers counted every day), on the fruit set following self- and open-pollination and on the effect of association of varieties in the orchards (choice, rate and placement of pollinisers).

Based on the results the rate of the overlap of the blooming times were calculated and varieties were assigned into five bloom time groups according to their main bloom. Self-fertility conditioned by natural self pollination was studied and good pollinisers were chosen (sweet, sour and duke cherry varieties) for the self-sterile and partially self-fertile varieties.

The necessity of bee pollination was proved by different pollination methods: natural self-pollination, artificial self-pollination, open pollination.

Keywords: fertility, flowering, pollination, sour cherry.

INTRODUCTION

Sour cherry production is an important industry and it is also an export item in Hungary. All except one of the registered and available 22 varieties are of Hungarian origin. The assortment is rather rich, moreover, regional selection and pedigree breeding promise currently new varieties.

Fertilisation in the *Pándy meggy*, then leading variety, has concerned the growers since the 19th century. Maliga (1942) surveyed the relevant literature. Self-fertility and its relation to fruit set obtained by open pollination was assessed in several Hungarian and foreign varieties by Nyéki (1974). All variants of *Pándy meggy* proved to be self-incompatible, and mutually inter-incompatible. Nyéki et al. (1992) called the

attention to the fact that the self-incompatible Pándy meggy cannot be fertilised by several self-fertile varieties (Debreceni bőtermő, Kántorjánosi, Újfehértói fürtös), i.e. they are inter-incompatible.

Sugar content of nectaries is high in sour cherry flowers; thus honeybees are attracted. Earlier observations stated the beneficial effect of bee activity in both self-incompatible as well as self-fertile varieties. i.e. fruit set increased substantially (Benedek et al. 1990).

In the present paper first of all the variability of fruit set depending on self- or alternatively open pollination in sour cherry varieties was analysed.

The objectives of the studies were to determine the blooming time, self and open

pollination of Hungarian sour cherry varieties.

METHODS

Observations of blooming dynamics followed the method applied by Nyéki (1974) on flowers distributed to the four quarters of the heavens on the end of branches, a sample comprising 100-500 flowers per variety. Starting with the beginning of bloom, every day at the same time (between 10 and 12 a.m.) the number of flowers according to their stage of development from the bud phase, opening and petal shedding was registered.

Flowers are considered to be open if the anthers and the pistil are easily recognised from above, and the stigma is green or yellow. The stigma of fading flowers started browning. Those dynamic observation served as a basis of determining the start of blooming (with the first open flowers), the main blooming period (when more than 50% of flowers opened) and the day of main bloom (when the number of open flowers reached the maximum) and the end of bloom (when the last flower faded).

According to the start of blooming varieties are grouped into 5 categories (early, medium early, intermediate, medium late, late).

Studies of fertilisation were done according to the method of Rudloff & Schanderl (1950).

Autogamy was investigated on 20-30 cm long branches of different orientation (N, E, S, W) covered still at bud stage with parchment paper bags. 5-10 isolated units comprising 100-400 flowers represented each variety. Paper bags were eliminated 3-4 days after the blooming period ended. The effect of artificial self pollination or geitonogamy was checked on isolated branches prepared as in the case of autogamy, except that the fully open flowers were hand pollinated once with

previously collected and stored (at +4°C) pollen of the same variety, subsequently, the bags were restored until the end of blooming period as in the case of autogamy.

Open pollinated flowers were observed, equally, on 5-10 branches 1.5-2 meter above ground level of four different orientations and comprising 100-500 flowers per variety.

Fruit set was registered and compared in all treatments at 1-2 weeks before fruit ripening.

RESULTS AND DISCUSSION

Blooming time

Sour cherry belongs to the group of medium early-blooming fruit trees but it is the latest among the stone fruits. At the start of blooming cultivated varieties cover a period of 4-7 days, during some seasons exceptionally also 10 days. Most authors distinguish three blooming time groups in sour cherry varieties. Present observations derived from three growing sites, 101 varieties, and a long period of years (1972-2002), which allowed the establishment of five blooming time groups. In Table 1 the propagated varieties are shown separately from the most important Hungarian and foreign varieties.

Neighbouring blooming time groups differ by 1-2 days only, depending on the season. In order to achieve considerable (more than 20%) fruit set a self-incompatible variety should be pollinated by a compatible polliniser, when their blooming period is overlapping each other by 70%, at least. In a system of 3 blooming time groups varieties belonging to the same group should be combined, whereas with 5 groups the overlap also of the neighbouring groups may prove to be sufficient for safe fruit set. Synchronous blooming or the extent of overlapping, however, may change yearly.

Variants of the variety *Pándy* differ in blooming time, thus *Pándy 48* is medium

Table 1

Blooming time groups of sour cherry varieties

Early	medium early	intermediate	medium late	late
Érdi bőtermő	Meteor korai Csengődi Cigánymeggy C. 404 Maliga emléke Pándy 48	Favorit Érdi nagygyümölcsű Korai pipacsmeggy	Érdi jubileum Cigánymeggy 7 Cigánymeggy 59 Pándy Bb. 119	Pándy 279 Debreceni bőtermő Kántorjánosi 3 Újfehértói fűrtös Oblacsinszka
Török meggy Egri fűrtös 102 Törpe meggy		Montmorency (Nagy Gobet) Cigánymeggy 3	Hartai meggy Fanal Cigánymeggy 215	Paraszt meggy Schattenmorelle (Latos meggy)

Table 2

Minimal and maximal overlap (%) of blooming periods in the *Pándy meggy* group of sour cherries with the potential polliniser varieties (Érd, 1972-1974)

♀	♂				
	Cigánymeggy 7	Cigánymeggy 59	Parasztmeggy	Schattenmorelle	Germersdorfi 57
Pándy 48	57-82	57-64	57-73	57-57	71-100
Pándy 279	75-100	75-90	80-90	75-80	60-90

early blooming, *Pándy 279* is late blooming. Blooming of *Pándy 48* is variable from year to year, thus the 70% overlap was hardly achieved with the medium late blooming *Cigánymeggy 7*, *Cigánymeggy 59* and late blooming *Parasztmeggy* and *Schattenmorelle* (Table 2). After having studied five potential pollinisers, only *Germersdorfi 57*, a sweet cherry variety proved to be a safe polliniser. *Pándy 279* seemed to overlap (by more than 70%) the bloom of *Cigánymeggy 7*, *Cigánymeggy 59*, *Parasztmeggy* and *Schattenmorelle*, whereas *Germersdorfi 57* was not sufficient every year.

Self-fertility

Self fertility of sour cherry varieties is a genetically determined property. In the extent of self-fertility there are substantial differences. In the growing practice, the self-incompatible varieties (e.g. types of the

Pándy meggy group), as unsafe yielders are losing space, whereas the popularity of self-fertile varieties is increasing.

Nyéki (1989) rated stone fruit varieties according to their extent of self-fertility into five groups. Sour cherry varieties observed on those three growing sites are assigned to those five groups as shown in Tables 3 and 4. In the assortment of sour cherries all variants between complete self-incompatibility and highly self-fertility are represented.

A high yield requires a 20-30% fruit set. Most of the sour cherry varieties are unable to set as much fruit at natural self pollination. Our data corroborate the statement of Rjabov & Rjabova (1970), which means that self-fertile sour cherry varieties are yielding safely every year because they are less subject to adversities due to weather and to growing site. Partially self-fertile varieties may produce 0-1% as well as more

Table 3

Comparison of fruit set obtained by open pollination
and by self-pollination in sour cherry varieties

Variety	Self-fertility groups	Self-pollination (%)	Open pollination (%)	Difference between self-and open pollination (%)
Cigánymeggy 1317 ¹	completely self-incompatible (0%)	0	19.5	19.5
Törpe meggy ¹		0	4.7	4.7
Mean		0	12.1	12.1
Cigánymeggy 60 ¹	self-incompatible (0.1-1 %)	0.2	6.1	5.9
Cigánymeggy 215 ¹		0.1	15.5	15.5
Török meggy ¹		0.3	6.4	6.4
Mean		0.2	9.3	9.1
Cigánymeggy C.404 ²	partially self-fertile (1.1-10 %)	6.3	28.8	22.5
Debreceni bőtermő ^{2,3}		5.5	22.8	17.3
Diemitzer ¹		7.1	20.2	13.1
Kántorjánosi ^{2,3}		5.0	20.1	15.1
Montmorency 1 ¹		5.3	32.5	27.2
Montmorency 2 ¹		2.0	17.5	15.5
Nagy Gobet ¹		1.2	9.5	9.5
Újfehértói fürtös ^{2,3}		6.0	23.1	17.1
Mean		4.8	21.8	17.0
Cigánymeggy 7 ¹	self-fertile (10.1-20%)	14.4	23.3	8.9
Cigánymeggy 59 ^{1,2}		13.6	23.9	10.3
Hartai meggy ^{1,2}		11.8	28.2	16.4
Latos meggy ¹		10.4	27.3	16.9
Montmorency 3 ¹		11.6	41.9	30.3
Parasztmeggy ^{1,2}		19.5	30.3	10.8
Schattenmorelle 2 ¹		18.5	38.9	20.4
Schattenmorelle 3 ¹		16.7	32.8	16.1
Schattenmorelle 4 ¹		16.9	28.3	11.4
Mean		14.5	29.0	14.5
Schattenmorelle 11	highly self-fertile (more than 20%)	25.3	29.4	4.1

Site and date of the observations: ¹ Érd-Elvira 1972-1974,
² Helvécia 1987-1988,
³ Újfehértó 1983-2002.

Table 4

Extreme values of fruit set obtained by self- and open pollination
in sour cherry varieties and types

Variety (type)	Self-incompatibility group	Self-pollination (%)			Open pollination (%)		
		minimum	maximum	difference	minimum	maximum	difference
Cigánymeggy 1317 ¹	completely self-incompatible (0.0%)	0.0	0.0	0.0	12.6	26.0	13.4
Törpe meggy ¹		0.0	0.0	0.0	2.6	6.7	4.1
Mean		0.0	0.0	0.0	7.6	16.4	8.8
Cigánymeggy 60 ¹	self-incompatible (0.1-1%)	0.0	0.6	0.6	0.0	8.9	8.9
Cigánymeggy 215 ¹		0.0	0.2	0.2	8.5	27.4	18.9
Török meggy ¹		0.0	0.9	0.9	1.4	12.7	11.3
Mean		0.0	0.6	0.6	3.3	16.3	13.0
Cigánymeggy C.404 ²	partially self-fertile (1.1-10%)	4.4	8.2	3.8	24.5	33.1	8.6
Debreceni bőtermő ^{2,3}		0.5	12.7	12.2	6.6	38.3	31.7
Diemitzer ¹		6.4	8.8	2.4	14.2	26.2	12.0
Kántorjánosi ^{2,3}		0.0	10.9	10.9	3.0	37.1	34.1
Montmorency 1 ¹		2.9	7.6	4.7	2.8	36.0	33.2
Montmorency 2 ¹		0.7	3.0	2.3	7.7	27.5	19.8
Nagy Gobet ¹		0.2	2.2	2.0	5.0	14.0	9.0
Újfehértói fűrtös ^{2,3}		0.7	18.7	18.0	3.0	42.5	39.5
Mean			2.0	9.0	7.0	8.4	31.8
Cigánymeggy 3 ¹	self-fertile (10.1-20%)	8.4	13.1	4.7	4.7	24.4	19.7
Cigánymeggy 59 ^{1,2}		8.5	26.7	18.2	14.4	22.5	8.1
Hartai meggy ^{1,2}		6.9	17.1	10.2	18.2	39.7	21.5
Latos meggy ¹		8.6	12.1	3.5	17.9	36.7	18.8
Montmorency 3 ¹		3.7	19.5	15.8	32.3	51.5	19.2
Paraszt meggy ^{1,2}		10.8	30.0	19.2	15.9	42.7	26.8
Schattenmorelle 2 ¹		12.8	29.0	16.2	27.9	46.5	18.6
Schattenmorelle 3 ¹		16.1	17.3	1.2	25.6	40.0	14.4
Schattenmorelle 4 ¹		9.6	24.2	14.6	27.5	29.1	1.6
Mean			8.8	21.6	12.8	19.3	35.2
Schattenmorelle 1 ¹	highly self-fertile (more than 20%)	15.7	33.5	17.8	19.0	39.7	20.7

Site and date of the observations: ¹Érd-Elvira 1972-1974,

²Helvécia 1987-1988,

³Újfehértó 1983-2002.

Table 5

Self- and open fertilisation of sour cherry varieties (%) (Újfehértó, 1983-2002)

Variety	Type of fertilisation	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	mean of years
Újfehértói fűrtös	self	18.7	11.8	2.7	10.0	3.2	4.2	2.4	9.2	6.3	5.7	2.6	3.2	1.3	6.9	-	1.7	0.7	1.4	5.0	13.0	5.8
	open	14.3	21.7	22.9	41.0	22.1	31.5	12.2	16.0	40.7	25.6	28.7	-	3.0	18.2	13.2	24.3	22.8	19.9	19.3	36.5	22.8
Kántorjánosi	self	9.1	7.9	1.8	4.2	3.9	3.9	6.3	6.0	40.4	8.2	4.1	5.1	0.0	4.4	-	1.7	5.5	1.8	4.1	10.9	4.9
	open	9.2	16.9	11.9	28.6	16.5	20.7	30.5	14.2	29.4	27.6	27.0	-	3.0	26.7	16.2	26.7	22.0	16.1	8.0	16.3	19.4
Debreceni bőtermő	self	8.5	2.2	2.2	5.4	5.4	6.3	12.7	5.5	10.6	6.1	3.6	4.8	8.3	2.9	-	4.0	5.0	0.5	3.9	11.2	5.7
	open	16.7	-	16.5	24.7	26.8	35.6	25.6	18.5	29.5	36.8	38.3	-	6.6	13.8	13.9	22.8	25.2	8.1	11.7	42.5	23.0

Table 6

Comparison of fruit set (%) in sour cherry varieties obtained by natural self-pollination (antogamy), artificial self-pollination (geitonogamy) and free pollination (Helyécia, 1988-1989)

Variety	1988			1989		
	Natural self-pollination (autogamy)	Artificial self-pollination (geitonogamy)	Open pollination	Natural self-pollination (autogamy)	Artificial self-pollination (geitonogamy)	Open pollination
Pándy 7	0	0	7.0	0	0	27.0
Debreceni bőtermő	1.4	17.2	9.4	6.8	21.8	31.8
Kántorjánosi	5.2	10.2	17.4	6.7	10.8	37.1
Cigánymeggy C 404	4.4	17.0	24.5	8.2	21.8	33.1
Újfehértói fűrtös	3.2	1.5	15.7	12.2	21.6	35.7
Cigánymeggy 59	8.6	28.7	31.0	11.7	32.6	36.6
Cigánymeggy 7	6.8	16.7	21.4	17.4	34.5	37.4
Parasztmeggy	14.7	23.1	33.6	17.1	39.7	42.7
mean	5.5	14.3	20.0	10.0	22.9	35.2

than 10% fruit set; variability of their yield is much higher than that of the self-fertile varieties (Table 4). Maximum fruit set of partially self-fertile varieties may amount 4.5 times their minimum set, whereas only 2.5 times in self-fertile varieties.

The high yearly variation of fruit sets is shown in Table 5. In the *Pándy meggy* type of varieties being partially self-fertile the yields show multiple differences depending on the season. Selfed flowers yield nearly 0% fruit, and the maximum yields did not reach 20% in any of them. As a mean of 19 years there were three varieties with similar rates of fruit set. Earlier observations (Nyéki, 1974) proved the beneficial effect of geitonogamy on the rate of fruit set in relation to natural self pollination or autogamy even in self-fertile varieties. In partially self-fertile varieties the advantage of geitonogamy was much less expressed than in the self-fertile group. However, natural autogamy allowed a higher variation in fruit set than did geitonogamy within the clone. Table 6 proves that with geitonogamy (hand pollination with pollen of the same clone) fruit set increased by a factor of 2-3 in relation to natural self-pollination or autogamy. Self-incompatible varieties (e.g. *Pándy 7*) did not set fruit even after geitonogamous hand pollination.

Open pollination

Fruit set on open pollinated flowers varied substantially according to growing site as well as to the season, which proves the high susceptibility of sour cherry to ecological adversities. Most decisive is the influence of weather on fruit set during the blooming period. Nyéki (1989) stated that fruit set of open pollinated *Pándy meggy* types is low and seasonally highly variable.

Tables 3 and 4 shows fruit sets of the varieties assigned to five self-fertility groups. Within the group of *Cigánymeggy* there are varieties, which are very poorly (e.g. *Cigánymeggy 60*), or well fertilised (e.g.

Hartai meggy, *Paraszt meggy*). Some favourable seasons may produce high (26-27%) fruit sets even in self- -incompatible varieties (e.g. *Cigánymeggy 215*, *Cigánymeggy 1317*). At present, only the best fertilised *Cigánymeggy* types are admitted to be propagated (*Cigánymeggy 7*, *Cigánymeggy 59*). Some varieties (e.g. *Montmorency 3*) may set fruit at rates more than 50%.

As was mentioned, self-incompatible and partially self-incompatible varieties yield more variably not only after self pollination but also with free pollination in relation to the self-fertile varieties.

Nyéki (1989) stated that the higher the self-fertility of a variety the higher yields are expected also with open pollination. A linear correlation exists between the rate of self-fertility and the productivity at open pollination at $P=0.01$ level. It is also proved by data appearing on Table 3. Completely and highly self-incompatible varieties set fruit with open pollination at rates of 12.1% and 9.3%, whereas self-fertile and highly self-fertile sour cherry varieties produced the highest fruit set: 29.0% and 29.4%.

CONCLUSIONS

1. Self-incompatible and partially self-incompatible sour cherry varieties must be planted with associated polliniser varieties. Appropriate polliniser varieties are to be chosen from groups of synchronous blooming period. If three blooming time groups are considered, the partners should belong to the same group, whereas from the five groups, they may be taken from the neighbouring blooming time groups too.
2. The rate of self- and open fertilisation is highly variable depending on the growing site as well as on the season. Variation of fruit set (and yield) is a good deal lower in self-fertile varieties than in self-incompatible and partially self-fer-

- tile varieties. Consequently, the safest yields are expected in self-fertile varieties.
3. Artificial self-pollination (geitonogamy) resulted in 2-3 times higher rates of fruit set in relation to natural self-pollination (autogamy), except in completely self-incompatible varieties. Highest fruit set is expected on open blooming flowers visited by bees. The advantage of open pollination is clearly evident in self-fertile and completely self-fertile varieties too. Consequently, it should be stated that bee pollination increased the yield not only in self-incompatible varieties but also in self-fertile sour cherry plantations.

REFERENCES

- Benedek P., Nyéki J., Szabó Z. (1990)
 - Cseresznye- és meggyfajták méhmegporzást befolyásoló tulajdonságai. *Kertgazdaság* 22(5):1-23.
- Maliga P. (1942) - Adatok a Pándy meggy virágbiológiájához. *Magyar Kir. Kertészeti Akad. Közl.* 8: 3-5.
- Nyéki J. (1974) - Meggyfajták virágzása és termékenyülése. Kandidátusi értekezés. *Kertészeti Egyetem*, Budapest (Kézirat).
- Nyéki J. (1989) - Csonthéjas gyümölcsűek virágzása, termékenyülése. *Akadémiai doktori értekezés.* MTA, Budapest (Kézirat).
- Nyéki J., Szabó Z., Salgim, S. S. (1992) - Self and cross compatibility of Hungarian sour cherry cultivars. *Acta Agr. Hung.* 41(1-2): 11-14.
- Rjabov, I. N., Rjabova, A. N. (1970) - Samoopylenie i perekrestnoe opylenie u cheresni, visni i vihsne-cheresni. *Trudii Gosud. Nikit. Botan.* Sada 4: 53-153.
- Rudloff C. F., Schanderl H. (1950) - Die Befruchtungsbiologie der Obstgewächse und ihre Anwendung in der Praxis. Stuttgart, Ulmer-Verlag.

BIOLOGIA KWITNIENIA I ZAPYLANIE WIŚNI (Prunus cerasus L.) ODMIAN WYSELEKCJONOWANYCH NA WĘGRZECH

Nyéki J., Szabó T., Szabó Z.

S t r e s z c z e n i e

Doświadczenia prowadzono w latach 1972 - 2002 w trzech miejscowościach: Erd, Helvecia i Ujfeherto, w których badano odmiany wiśni uprawiane w różnych zestawach odmianowych. Obserwacje obejmowały: biologię kwitnienia (czas rozpoczęcia, okres trwania, największe nasilenie), dynamikę kwitnienia (liczba otwieranych codziennie kwiatów) oraz zawiązywanie owoców i wpływ na to sąsiedztwa odmian znajdujących się w sadzie.

Opierając się na uzyskanych wynikach oceniono stopień nakładania się okresów kwitnienia różnych odmian, a na podstawie okresu największego nasilenia kwitnienia wszystkie odmiany podzielono na 5 grup. Badano samopłodność i zdolność do naturalnego samozapyłania się oraz wybrano dobre zapylacze dla odmian samosterylnych i częściowo samosterylnych.

Potrzebę zapylenia przez pszczoły oceniono stosując różne sposoby zapyłania; naturalne samozapylenie, sztuczne samozapylenie i wolne zapyłanie.

Słowa kluczowe: zapłodnienie, kwitnienie, zapyłanie, wiśnia.